

# The Effectiveness of Psychoeducational Group Intervention on Post-Traumatic Growth: A Randomized Clinical Trial

Negin Masoum Abadi<sup>1</sup>, Jamileh Zarei<sup>2\*</sup>, Seyede Saleheh Mortazavi<sup>3,4</sup> and Ramesh Omranipour<sup>5</sup>

1. Tabriz University of Medical Sciences, Tabriz, Iran

2. Department of Health Psychology, Research Center of Spiritual Health, School of Behavioral Sciences and Mental Health, Iran University of Medical Sciences, Tehran, Iran

3. School of Behavioral Sciences and Mental Health, Tehran Institute of Psychiatry, Iran University of Medical Sciences (IUMS), Tehran, Iran

4. Spiritual Health Research Center, Iran University of Medical Sciences, Tehran, Iran

5. Breast Disease Research Center, Tehran University of Medical Sciences, Tehran, Iran

## \* Corresponding author

### Jamileh Zarei, MD

Department of Health Psychology,  
Research Center of Spiritual Health,  
School of Behavioral Sciences and  
Mental Health, Iran University of Medical  
Sciences, Tehran, Iran

**Email:** jamileh.zarei@gmail.com

**Received:** 1 Apr 2019

**Accepted:** 23 Jun 2019

### Citation to this article:

Masoum Abadi N, Zarei J, Mortazavi SS, Omranipour R. The Effectiveness of Psychoeducational Group Intervention on Post-Traumatic Growth: A Randomized Clinical Trial. *J Iran Med Council.* 2019;2(6):174-183

## Abstract

**Background:** Breast cancer is a chronic disease that leads to many psychological problems that are involved in treatment and adherence to treatment. The aim of this study was to evaluate the effect of psychoeducational group intervention on psychological factors that help the patients in the treatment process.

**Methods:** In this study, 80 patients were selected by using convenience sampling at Imam Khomeini Hospital. They were randomly allocated either to the intervention group or to the control group. All participants completed the questionnaires of post-traumatic growth, quality of life, subjective well-being, adherence to treatment and social support. The results were analyzed by SPSS-18 software using univariate and multivariate analysis.

**Results:** According to the results, the applied intervention had a significant effect on post-traumatic growth as a primary outcome, and also on subjective well-being and adherence to treatment as secondary outcomes.

**Conclusion:** Psychoeducational group intervention in women with breast cancer can affect the treatment and provide better response to treatment through the improvement of post-traumatic growth, subjective well-being and adherence to treatment.

**Keywords:** Breast cancer, Educational intervention, Post traumatic growth, Randomized clinical trial

## Introduction

Cancer is one of the major causes of death in the world and the third cause of death in Iran (1). Common complications of cancer, such as fear, loneliness, death anxiety, alteration in relationships, tolerance of painful medical procedures, and uncertain future may lead to distress in cancer patients (1). Breast cancer, the second most common cancer in the world (2), accounts for 25.1% of all cancers with a prevalence of 1.67 million cases worldwide (3). In Iran, breast cancer is the first diagnosis among all cancers and its incidence rate is higher than other cancers (4).

The survival rate of breast cancer has increased significantly over the past 20 years. Though most breast cancer patients face a variety of problems after treatment, they return to daily life with appropriate psychosocial interventions (2,5). However, the response to the disease may vary in some individuals and some cases require a catalyst to achieve higher levels of psychological growth. This is called post-traumatic growth, and in recent years, researchers have gradually shifted their focus from traumatic event to positive outcomes.

The occurrence of incurable illness requires the reinforcement of positive psychological aspects of individuals and strengthening their skills in dealing with traumatic events throughout their lives. Such problems may not destroy the whole life of individuals; quite contrary, these stressful events, along with negative consequences, may bring positive consequences for their individual and social lives (6).

Several studies have investigated the necessity of psychological interventions for breast cancer patients and the role of these interventions in reducing the effect of illness on well-being and quality of life as well as facilitating post-traumatic growth. Group support, individual counseling, stress management, and psychological training are examples of psychological interventions that have been shown to have significant effects on these variables in women with breast cancer (7).

Subjective well-being is an important prognostic factor for breast cancer adherence to treatment. It refers to individuals' assessment and perception of their quality of life, including the quality of their social, psychological, and emotional actions. In other

words, subjective well-being implies how individuals evaluate their lives, including variables such as life satisfaction, marital satisfaction, positive moods, lack of depression and anxiety (8).

Social support, in particular, facilitates recovery, enhances access to medical treatment, and enhances the overall quality of life (9). A sense of social support in the face of uncertainty about life events reinforces a sense of self-control (10). Social support is considered an active stress management strategy that may reduce stress levels (11). Some other studies have shown that access to social support during and after a traumatic event can reduce levels of depression, anxiety, and other physical and mental disorders among those exposed to some traumatic events (12). In relation to post-traumatic growth, social support is seen as a direct predictor of positive change after unpleasant experiences (13).

Based on existing research literature such as the studies of Shorey *et al* (14) and Brand *et al* (15), psychoeducation can increase social support. Social support can also increase post-traumatic growth, according to Scignaro *et al* (16), Schroevers *et al* (17), and Bozo *et al* (18). Therefore, in this study, the role of social support as a mediator variable was examined. Since there is no comprehensive study about any positive psychological feature that affects treatment of cancer, this study aimed to assess the effectiveness of psychoeducational group intervention on post-traumatic growth, quality of life, subjective well-being, and adherence to treatment for women with breast cancer by examining the mediating role of social support.

## Materials and Methods

### Study design

The present study was a randomized controlled trial. Post-traumatic growth was the primary outcome and quality of life, subjective well-being, and adherence to treatment were secondary outcomes. In addition, the mediating role of social support variable was examined.

### Participants

Eighty women with breast cancer at stages one and two in the 1<sup>st</sup> and 3<sup>rd</sup> wards of Cancer Institute at Imam Khomeini Hospital were selected after obtaining informed consent. The sampling method was convenience sampling.

Then, they were allocated to the intervention and control groups using block randomisation. Figure 1 shows the initial number of cases and the cause of losing the individuals in the final sample.

Inclusion criteria were being at least 20 years of age, cancer diagnosis by breast cancer specialist at stage 1 and 2, minimum basic literacy and willingness, and consent to participate in the study. Exclusion criteria included less than two weeks duration of illness, history of known psychiatric disorder, clinical symptoms of anxiety and depression, incomplete answering to questions, participation in some psychological intervention to alleviate the psychological symptoms associated with the disease, having a co-morbid condition or cognitive disorder.

**Measurements**

First, socio-demographic measures were recorded (age, education, employment, marriage status, kind of therapy, and stage of cancer). Patients completed validated Iranian versions of the following

questionnaires. The Posttraumatic Growth Inventory (PTGI) developed by Tedeschi and Calhoun was used to investigate the concept of post-traumatic growth. The tool has 21 items that evaluate five areas of psychological development after facing a stressful trauma (new situations, relationships with others, appreciation and appreciation for life, personality strengths, and spiritual change) (19). In Iran, the obtained Cronbach’s alpha for this inventory was 0.92 and its reliability was 0.94 (1).

Quality of life was measured using the Quality of Life Questionnaire (EROTC QLQ-C30). This questionnaire belongs to the European Organisation for Research and Treatment of Cancer to assess the quality of life in cancer patients. It measures nine areas of symptoms (Fatigue, nausea and vomiting, pain, shortness of breath, sleep deprivation, loss of appetite, constipation, diarrhea, and financial problems) and one general area of quality of life (20-23). The validity and reliability of this questionnaire have been evaluated by Marashi *et al* (22). The patients also completed Breas




Figure 1. The initial samples, causes of sample loss.

Cancer-Specific Quality of Life Questionnaire (QLQ-BR23). The questionnaire consists of 23 questions measuring four functional domains (sexual function, sexual pleasure, and body image and prospects) and four symptom domains (hair loss discomfort, treatment side effects, arm symptoms, and breast symptoms). The validity and reliability of this questionnaire in Iran have been assessed by Shobeiri F *et al* (24).

Patients were also assessed by Subjective Well-being Scale (SWS) designed by Keyes CL (8). This scale includes three dimensions of emotional well-being (12 questions), psychological well-being (18 questions) and social well-being (15 questions). Adherence was assessed using the 8-item Morisky Medication Adherence Scale (MMAS-8). The range of average scores is zero to 8. The score of less than six is classified as poor compliance, 6 to 8 is moderate and 8 is appropriate compliance. Validation of the Persian version of this tool in Iranian patients with hypertension

was performed by Moharamzad Y *et al* (25).

### **Intervention**

The present study utilized structured psychoeducational intervention for cancer patients developed by Fawzy FI and Fawzy NW (26) to investigate the effectiveness of group psychological training intervention on post-traumatic growth, quality of life, mental well-being, and evaluation. Table 1 presents the contents of this protocol. The intervention group received six sessions of group training for 90 minutes twice a week by a psychotherapist trained in the field of patient education. The control group received common treatments including chemotherapy, surgery and medical consultation.

### **Ethics approval**

At the beginning and before the actual implementation of the research, the purpose of the study and the method of training for both the intervention and control groups

**Table 1.** The content of psychological training sessions

Agenda	Sessions
Referrals and explanation of the structure of the meetings	1
<b>Health Education</b>	2
Health care information ,including types of cancer Different treatment methods (Lumpectomy, mastectomy, radiation therapy, chemotherapy, and tamoxifen) Good follow-up routines (Monthly self-examination of breast, annual mammography, and annual medical examinations) Group discussion	
<b>Stress management</b>	3
Identifying sources of stress Personal reactions to stress (Physiological, psychological and behavioral reactions) Problem-solving Change of attitude Group discussion	
<b>Relaxation training</b>	4
Explanation of relaxation exercises Progressive muscle relaxation followed by guided imaging Group discussion	
<b>Coping skills</b>	5
Active behaviors Active cognitive methods Avoidance methods Coping scenarios (with picture) group discussion	
Summary Group discussion	6

were explained, and their consent to participate in the study was obtained. This study was approved as a minimal-risk research by the Ethics Committee of Iran University of Medical Sciences (Approval reference number: IR.IUMS.REC.1397.276).

## Results

The mean age of the intervention group was  $48 \pm 9$ , and in the control group was  $51 \pm 8$ . The mean duration of disease in the intervention group was five months and in the control group was six months. Table 2 shows the baseline characteristics of the two study groups.

In the univariate analysis, as shown in table 3, changes in outcome variables were significantly different between the study groups in the variables of post-traumatic growth, subjective well-being and adherence to treatment.

To evaluate the mediating effect of social support and potential confounding variables (demographic variables), the crude effect of intervention variable in a regression model with only outcome variables (post-traumatic growth, subjective well-being, quality of life and treatment adherence) and the main independent variable (intervention) was calculated.

**Table 2.** Baseline characteristics of intervention and control group

Variable	Intervention		Control		
	Frequency	Percentage	Frequency	Percentage	
Education	Primary school	9	30	5	16.7
	Secondary school	18	26.7	12	40
	Undergraduate	2	6.7	3	10
	Graduate	1	3.3	0	0
Employment	Employed	4	13.3	7	23.3
	Non-employed	26	86.7	23	76.7
Marriage status	Married	25	83.3	28	93.3
	Single	5	16.7	2	6.7
Kind of therapy	Mastectomy	14	46.7	11	36.7
	Lumpectomy	16	53.3	19	63.3
Stage of cancer	1	12	40	15	50
	2	18	60	15	50

**Table 3.** Comparison of changes in outcome variables between two study groups in univariate analysis

Outcome variables	Group	N	Average change after - before	SD	p-value
Post-traumatic growth change	Intervention	30	21.56	14.00	<0.001
	Control	30	- 7.20	14.44	
Subjective well-being change	Intervention	30	32.46	16.84	<0.001
	Control	30	7.70	31.36	
Quality of life change	Intervention	30	- 13.96	9.953	0.24
	Control	30	- 9.46	16.78	
Adherence change	Intervention	30	1.16	2.03	0.01
	Control	30	- 0.13	1.94	

**Table 4.** Assessing the mediator effect of social support and potential confounding factor of age, education, and occupation (Multivariate analysis)

	Post-traumatic growth	Well-being	Adherence
Crud effect size	29.06	24.76	1.30
Social support	29.11	24.78	1.30
Age	29.60	23.22	1.18
Education	1.29	24.69	29.09
Occupation	1.38	23.12	28.32

The effect size was 29.06, 24.76, 4.49 and 1.3 respectively. The p-value was less than 0.001 in post-traumatic growth and subjective well-being 0.01, for adherence and was not statistically significant for the quality of life (p-value=0.24)

Then, the effect size mentioned above for the post-traumatic growth, subjective well-being, and treatment adherence variables was recalculated in the presence of social support variables and potential confounding variables in separate models (adjusted model). The size of the primary crude effect was compared with the effect size calculated in the secondary model. Table 4 shows the result of multivariate analysis. Due to the difference between the crude effect size and the adjusted as less than 10%, there is not a mediator and potential confounder variable.

## Discussion

Based on the result, psychological training has been effective in improving post-traumatic growth. The results are in line with the findings of previous literature (27-30). Garlan *et al* (31) showed that participants in the intervention group reported more traumatic growth than the control group.

In explaining the effectiveness of psychological training on post-traumatic growth in breast cancer patients, as Lelorain and Bonnaud-Antignac (32) have pointed out, talking to others can stimulate reassessment, facilitate the reconsideration of world assumptions, create meaning, and reinforce emotion expression. Talking and thinking about loss has been associated with finding meaning and post-traumatic growth in other stress-stricken populations. According to the ontological and cognitive processing models, the perceived threat from breast cancer is positively related to traumatic growth (2).

What is important in post-traumatic growth in cancer patients is that they see the stress associated with the disease as a challenge for personal growth, understanding the meaning of life, reviewing their lives, and their connection to the world (32). Lelorain *et al* showed that positive, active, relational, religious, and partially consistent coping has a significant effect on post-traumatic growth. Coping strategies play an important role in traumatic growth. In many cancer situations, adaptive coping such as religious coping, acceptance, or reinterpretation, has been positively

associated with post-traumatic growth, whereas maladaptive coping such as substance use or self-blame is not associated with growth (32).

The perception of the stressful event threatens one's core belief system. The core beliefs are "A general set of one's beliefs about the world, how one functions and one's place in it." Following a stressful event, one's core beliefs may be challenged, and the person's hypothetical world may become unstable, uncontrollable, and unpredictable. To manage distress (as well as challenge towards core beliefs) and reevaluate pre-traumatic cognitive schemas, survivors use various cognitive mechanisms such as coping strategies, social support seeking, cognitive participation (Uninvolved thinking, and meditation). They begin by trying to give meaning to a stressful experience. Group psychosocial interventions may enhance post-traumatic growth by enhancing emotional expression, cognitive processing, and positive evaluation of a stressful event (32).

Although previous research suggested that psychoeducation had a positive effect on patients' quality of life (33,34), no such results were found in this study. This may be due to participation of individuals in the post-test which was performed after the surgery, with patients suffering from a variety of physical problems such as pain, and post-operative complications such as fatigue, nausea, and vomiting. Another reason may be related to short-term education because studies that showed an improvement in quality of life after intervention provided long-term education for patients. Long-term interventions are needed to evaluate the effectiveness of other types of psychosocial support in improving the quality of life in early-stage breast cancer patients. Regarding the third hypothesis of the research suggesting that group training in psychological training is effective on increasing subjective well-being in patients with breast cancer, the findings of the study showed that group psychological training had a significant effect on promoting the mental well-being of women with breast cancer. This is in line with the results of the studies by Al-Sulaiman *et al* (7) and Dow Meneses K *et al* (34).

In explaining the effectiveness of psychological education on improving the well-being of patients with breast cancer (35,36), one's psychological resources

influence one's ability to use social support to improve subjective well-being. In the context of the group, supportiveness can come from group members. Group support for cancer patients is a source of emotional support. And there is evidence that for isolated individuals, the group can play such a role.

Psychosocial interventions can improve patients' knowledge and awareness of emotional and physical responses to stress and increase their confidence, help manage pain and other physical and emotional symptoms; they can help to reduce uncertainty, stress, anxiety, helplessness and fear of unknown conditions and psychological discomfort through relaxation training and other techniques which consequently improve psychological and physiological adaptation indexes. Health-promoting behaviors such as exercise, diet, sleep, and medication psychosocial skills and the use of interpersonal skills to build and maintain social support, coping strategies in helping to identify and change negative beliefs, beliefs and expectations, increasing ability to control thoughts, beliefs, and attitudes, as well as more positive attitudes to clinical trials all enhance the psychological well-being of patients with cancer.

Concerning the fourth hypothesis of research suggesting that group training in psychological training is effective in increasing adherence to treatment for patients with breast cancer, the findings of the study showed that group psychological training has significant effect on promoting adherence to treatment of women with breast cancer and this is consistent with the results of studies by Jeffries SA *et al*, (37) and Myers RE *et al* (38).

In explaining the effectiveness of psychological training on promoting adherence to treatment for patients with breast cancer, it can be said that according to Myers RE *et al* (38), forgetfulness, anxiety, lack of knowledge and skills, and misconceptions about therapeutic measures can be improved. Interventions in adherence to treatment include patient education strategies, including education about chronic diseases such as cancer, the benefits of treatment, and the serious consequences of non-compliance. And behavioral adjustment strategies include awareness of the benefits of follow-up and the high costs incurred by non-compliance (39). Training these topics can increase adherence to treatment for cancer patients.

Also, the results of this study are inconsistent with the findings of Al- Sulaiman *et al* (7). In this study, crisis counseling and psychological education did not have a significant effect on adherence to treatment for patients with breast cancer. This may be due to differences in mean age of the groups, instruments used, and duration of intervention.

The fifth hypothesis of the research about the role of group intervention in psychological education and development of post-traumatic social support was not confirmed in this study. Although studies have shown the mediating role of social support, according to the studies of Shorey S *et al* (14) and Brand EF *et al* (15), psychological education can increase social support. And social support can also increase post-traumatic growth, according to studies by Cobb (10), Scignaro M *et al* (16), Schroevers MJ *et al* (17), and Bozo O *et al* (18). Intermediary role failed because there are other variables between psychological education and social support or between social support and traumatic growth. The patients' conditions in the ward and during the surgery and the short duration of the intervention may also play a role, which requires further research.

The evidence from this study is also consistent with the sixth hypothesis, in line with the researches of Garlan RW *et al* (31), Rolnick SJ *et al* (39), and Baglama B *et al* (40), suggesting that age, education, occupation and marital status of individuals do not have much effect on the benefits of psychological training. While in marriages belonging to an ethnic minority group, higher education, employment, and youth are positively correlated with post-traumatic growth, the results suggest that these socio-demographic variables account for diversity (41). Few have been studied in our sample. In other words, the response to cancer is not determined so much by the concrete properties of the situation and the individual, but by the subjective reactions to the experience (*e.g.* disability, controllability, a threat to life, etc.) and the perceived interpersonal context (42).

### **Research limitations**

There are some limitations to this study. Short-term intervention due to limited hospitalization of patients inward is the first limitation. Moreover, due to the nature of the intervention blindness is impossible for participants and the researcher. But the person who

analyzed the data did not know the cases.

## Conclusion

Psychological interventions, such as psychoeducation,

have positive effects on traumatic growth, subjective well-being, and adherence to treatment. However, no positive results were found in this study regarding the

---

impact of psychological education on quality of life.

## References

1. Heidarzadeh M, Rassouli M, Brant JM, Mohammadi-Shahbolaghi F, Alavi-Majd H. Dimensions of Posttraumatic Growth in Patients with Cancer: A Mixed-Method Study. *Cancer Nurs* 2018;41(6):441-9.
2. Norouzi H, Rahimian-Boogar I, Talepasand S. Effectiveness of mindfulness-based cognitive therapy on posttraumatic growth, self-management, and functional disability among patients with breast cancer. *Nursing Practice Today* 2017;4(4):190-202.
3. Barthakur MS, Sharma MP, Chaturvedi SK, Manjunath SK. Posttraumatic growth in women survivors of breast cancer. *Indian J Palliative Care* 2016;22(2):157-62.
4. Akbari M, Lotfi Kashani F, Vaziri S. The efficacy of four-factor psychotherapy on increasing sexual self-esteem in breast cancer survivors. *Iranian J of Breast Disease*. 2017;10(1):48-60.
5. Lim SM, Kim HC, Lee S. Psychosocial impact of cancer patients on their family members. *Cancer Res Treat* 2013;45(3):226.
6. Wu X, Kaminga AC, Dai W, Deng J, Wang Z, Pan X, et al. The prevalence of moderate-to-high posttraumatic growth: A systematic review and meta-analysis. *J Affect Disord* 2019 Jan 15;243:408-15.
7. Al-Sulaiman RJ, Bener A, Doodson L, Al Bader SB, Ghuloum S, Lemaux A, et al. Exploring the effectiveness of crisis counseling and psychoeducation in relation to improving mental well-being, quality of life and treatment compliance of breast cancer patients in Qatar. *Int J Womens Health* 2018;10:285-96.
8. Keyes CL, Shapiro AD. Social well-being in the United States: A descriptive epidemiology. How healthy are we? In: Brim OG, Ryff CD, Kessler RC, (Eds.). *The John D. and Catherine T. MacArthur foundation series on mental health and development. Studies on successful midlife development. How healthy are we?: A national study of well-being at midlife.* USA: University of Chicago Press; 2004.P.350-72.
9. Rizalar S, Ozbas A, Akyolcu N, Gungor B. Effect of perceived social support on psychosocial adjustment of Turkish patients with breast cancer. *Asian Pac J Cancer Prev* 2014;15(8):3429-34.
10. Cobb S. Social support as a moderator of life stress. *Psychosomatic Medicine* 1976;38(5):300-14.
11. Flannery Jr RB. Social support and psychological trauma: A methodological review. *J Traumatic Stress* 1990;3(4):593-611.
12. Regehr C, Hill J, Knott T, Sault B. Social support, self-efficacy and trauma in new recruits and experienced firefighters. *Stress and Health* 2003;19(4):189-93.
13. Prati G, Pietrantonio L. Optimism, social support, and coping strategies as factors contributing to posttraumatic growth: A meta-analysis. *J Loss and Trauma* 2009;14(5):364-88.
14. Shorey S, Chan SWC, Chong YS, He HG. A randomized controlled trial of the effectiveness of a postnatal psychoeducation program on self-efficacy, social support and postnatal depression among primiparas. *J Adv Nurs* 2015;71(6):1260-73.
15. Brand EF, Lakey B, Berman S. A preventive, psychoeducational approach to increase perceived social support. *Am J Community Psychol* 1995;23(1):117-35.
16. Scrignaro M, Barni S, Magrin ME. The combined contribution of social support and coping strategies in predicting

post-traumatic growth: a longitudinal study on cancer patients. *Psycho-Oncology* 2011;20(8):823-31.

17. Schroevers MJ, Helgeson VS, Sanderman R, Ranchor AV. Type of social support matters for prediction of posttraumatic growth among cancer survivors. *Psycho-Oncology* 2010;19(1):46-53.

18. Bozo Ö, Gündoğdu E, Büyükaşık-Çolak C. The moderating role of different sources of perceived social support on the dispositional optimism-posttraumatic growth relationship in postoperative breast cancer patients. *J Health Psychol* 2009;14(7):1009-20.

19. Tedeschi RG, Calhoun LG. The posttraumatic growth inventory: Measuring the positive legacy of trauma. *J Trauma Stress* 1996;9(3):455-71.

20. Matsuda A, Yamaoka K, Tango T, Matsuda T, Nishimoto H. Effectiveness of psychoeducational support on quality of life in early-stage breast cancer patients: a systematic review and meta-analysis of randomized controlled trials. *Qual Life Res* 2014;23(1):21-30.

21. Diener E. Subjective well-being. *Psychological Bulletin* 1984;95(3):542-75.

22. Marashi T, Taherianfar Z, Etemad K, Rakhsha A. [Survey of quality of life of patients with breast cancer among women at Shahid Beheshti University of Medical science and Health service 2017]. *Journal Health in the Field* 2018;6(1):1-8. Persian.

23. Esfandiari B, Kazemi M, Amanpour E, Tizfahm T. The relationship between religion, spiritual, health, hope, and quality of life in cancer patients. *BCCR* 2014;6(4):28-36.

24. Shobeiri F, Nikravesht A, Masoumi SZ, Heidari Moghaddam R, Karami M, Badafreh M. [The effect of exercise counseling on functional measures of quality of life in women with breast cancer]. *J Education and Community Health* 2015;2(1):1-9. Persian.

25. Moharamzad Y, Saadat H, Shahraki BN, Rai A, Saadat Z, Aerab-Sheibani H, et al. Validation of the Persian version of the 8-item Morisky Medication Adherence Scale (MMAS-8) in Iranian hypertensive patients. *Glob J Health Sci* 2015;7(4):173.

26. Fawzy FI, Fawzy NW. A structured psychoeducational intervention for cancer patients. *Gen Hosp Psychiatry* 1994 May;16(3):149-92.

27. Aggarwal B, Liao M, Allegrante JP, Mosca L. Low social support level is associated with non-adherence to diet at one year in the Family Intervention Trial for Heart Health (FIT Heart). *J Nutrition Education and Behavior* 2010;42(6):380-8.

28. Shorey S, Chan SWC, Chong YS, He HG. A randomized controlled trial of the effectiveness of a postnatal psychoeducation program on self-efficacy, social support and postnatal depression among primiparas. *J Adv Nurs* 2015;71(6):1260-73.

29. Hashemian K, Pourshahriari MS, Bani Jamali MAS, Golestanie Bakht T. Investigating the relationship between demographic characteristics and subjective biodiversity and happiness in Tehran population. *J Educational Psychology Studies* 2007;3(3):139-63.

30. Pat-Horenczyk R, Perry S, Hamama-Raz Y, Ziv Y, Schramm-Yavin S, Stemmer SM. Posttraumatic growth in breast cancer survivors: Constructive and illusory aspects. *J Trauma Stress* 2015;28(3):214-22.

31. Garlan RW, Butler LD, Siegel ERA, Spiegel D. Perceived benefits and psychosocial outcomes of a brief existential family intervention for cancer patients/survivors. *Omega-Journal of Death and Dying* 2011;62(3):243-68.

32. Lelorain S, Bonnaud-Antignac A, Florin A. Long term posttraumatic growth after breast cancer: prevalence, predictors, and relationships with psychological health. *J Clin Psychol Med Settings* 2010;17(1):14-22.

33. Matsuda A, Yamaoka K, Tango T, Matsuda T, Nishimoto H. Effectiveness of psychoeducational support on quality of life in early-stage breast cancer patients: a systematic review and meta-analysis of randomized controlled trials. *Qual Life Res* 2014;23(1):21-30.

34. Dow Meneses K, McNees P, Loerzel VW, Su X, Zhang Y, Hassey LA, editors. Transition from treatment to survivorship: effects of a psychoeducational intervention on quality of life in breast cancer survivors. *Oncol Nurs Forum* 2007;34(5):1007-16.
35. Ram S, Narayanasamy R, Barua A. Effectiveness of group psycho-education on well-being and depression among breast cancer survivors of Melaka, Malaysia. *Indian J Palliat Care* 2013;19(1):34.
36. Bloom JR, Stewart SL, Johnston M, Banks P, Fobair P. Sources of support and the physical and mental well-being of young women with breast cancer. *Soc Sci Med* 2001;53(11):1513-24.
37. Jeffries SA, Robinson JW, Craighead PS, Keats MR. An effective group psychoeducational intervention for improving compliance with vaginal dilation: a randomized controlled trial. *Int J Radiat Oncol Biol Phys* 2006;65(2):404-11.
38. Myers RE, Chodak GW, Wolf TA, Burgh DY, McGrory GT, Marcus SM, et al. Adherence by African American men to prostate cancer education and early detection. *Cancer* 1999;86(1):88-104.
39. Rolnick SJ, Pawloski PA, Hedblom BD, Asche SE, Bruzek RJ. Patient characteristics associated with medication adherence. *Clin Med Res* 2013;11(2):54-65.
40. Baglama B, Atak IE. Posttraumatic growth and related factors among postoperative breast cancer patients. *Procedia-Social and Behavioral Sciences* 2015;190:448-54.
41. Sears SR, Stanton AL, Danoff-Burg S. The yellow brick road and the emerald city: benefit finding, positive reappraisal coping, and posttraumatic growth in women with early-stage breast cancer. *Health Psychol* 2003;22(5):487-97.
42. Ginzburg K, Ein-Dor T. Posttraumatic stress syndromes and health-related quality of life following myocardial infarction: 8-year follow-up. *Gen Hosp Psychiatry* 2011;33(6):565-71.