

# Association between Sleep Duration and Quality with Risk of Gestational Diabetes Mellitus

Shirin Niroomanesh<sup>1</sup>, Mona Mohseni<sup>2</sup>, Maryam Gholizadeh<sup>2</sup>, Nasim Shokuhi<sup>2</sup> and Nafiseh Saedi<sup>2\*</sup>

1. Maternal, Fetal, and Neonatal Research Center, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Obstetrics and Gynecology, Yas Hospital, Tehran University of Medical Sciences, Tehran, Iran

## Abstract

**Background:** Several studies have been conducted on the relationship between sleep disorders and gestational diabetes. In this study, the relationship between the quality and quantity of sleep and the incidence of gestational diabetes in pregnant women was assessed.

**Methods:** In this case-control study, the quality and quantity of sleep in pregnant women attended to the Perinatology clinic at Yas Hospital, Tehran, Iran between June 2016 to September 2017 and its relationship with gestational diabetes mellitus was assessed. In this research, 3000 pregnant women at 24 to 28 weeks of pregnancy were enrolled. Gestational diabetes screening test was initially performed on Glucose Challenge Test (GCT) with administration of 50 gr of glucose. For abnormal cases, Oral Glucose Tolerance Test (OGTT) with administration of 100 gr of glucose after an overnight fasting (8-10 hours) was requested. According to the Gestational Diabetes Mellitus (GDM) diagnosis criteria based on the American Diabetes Association (ADA), women were divided into two groups of case and control (290 diabetic cases and 2710 non-diabetic cases). The Pittsburgh Sleep Quality Index (PSQI) and the participants' demographic information were collected.

**Results:** Ultimately, 290 of 3,000 women in the study, based on the ADA criteria were diabetic and non-diabetic patients were 2710 women. Sleep quality in the diabetic mothers was significantly worse than non-diabetic mothers and the duration of sleep of them was higher than non-diabetic mothers ( $p < 0.05$ ).

**Conclusion:** Quality and quantity of sleep can be associated with the incidence of gestational diabetes and diabetic mothers have worse sleep quality and longer sleep times than non-diabetic mothers.

**Keywords:** Gestational diabetes mellitus, Pregnancy, Sleep disorder

## \* Corresponding author

**Nafiseh Saedi, MD**

Department of Obstetrics and Gynecology, Yas Hospital, Tehran University of Medical Sciences, Tehran, Iran

**Email:** nsaedi@sina.tums.ac.ir

**Received:** 10 Nov 2019

**Accepted:** 25 Dec 2019

## Citation to this article:

Niroomanesh Sh, Mohseni M, Gholizadeh M, Shokuhi N, Saedi N. Association between sleep duration and quality with risk of gestational diabetes mellitus. *J Iran Med Council.* 2020;3(1):23-28.

## Introduction

Gestational Diabetes Mellitus (GDM) that has shown an increasing trend worldwide is accompanied with adverse short-term and long-term outcomes for both mother and fetus including cardiovascular disorders, diabetes mellitus type II, childhood obesity, *etc* (1,2). GDM is multi-factorial and recognition of contributing factors would help better programming in decreasing the incidence rate of disease and possible outcomes (2). Sleep disorders especially elongated sleep, poor-quality sleep, and breathing disorders during sleep are common in pregnancy with possible etiological role of hormonal alterations, anxiety, and somatic problems (3,4). In previous studies, the association between sleep disorders and gestational adverse outcomes such as preeclampsia, elongated labor duration, increased cesarean rate, Intra Uterine Growth Retardation (IUGR), and preterm labor has been demonstrated (3-5).

Also, some studies were conducted on the association between sleep disorders in non-pregnant women and risk of glucose metabolism disorders, insulin resistance, and diabetes type II (6-8). Sleep is an important human requirement and the sleep quality in high-risk pregnant women with mild pre-eclampsia and GDM decreases (8,9). The association between sleep-related respiratory disorders and risk of diabetes mellitus has been shown (9) but the association with GDM and also the role of duration and quality of sleep is not yet clear. Clarifying such association may help to determine the role of contributing factors in GDM which are useful in decreasing its incidence rate (10-15). Hence, in this study, the association between GDM with duration and quality of sleep was assessed.

## Materials and Methods

In this case-control study, 3040 pregnant women at 24 to 28 gestational weeks were assessed who were seeking obstetric cares from perinatology clinic of Yas Hospital, Tehran, Iran from June 2016 to September 2017. The study was approved by local ethical committee with recorded code of 9311290021. This study was performed in accordance with Helsinki Declaration and the informed consent was signed by all participants.

The pregnant women with the history of type 1, 2

diabetes mellitus, history of respiratory syndrome, history of chronic heart or renal disease (n=40) were excluded, then Glucose Challenge Test (GCT) (blood glucose 1 hour post-50-gr glucose) was measured for 3000 pregnant women. All tests were done in a single lab of Yas Hospital. In the cases with abnormal blood glucose levels (GCT  $\geq 135$ ), OGTT with 100 gr glucose after 8-10 hours of overnight fasting was requested and blood glucose levels were measured before and 1, 2 and 3 hours after administration of 100 gr of glucose. The GDM was diagnosed according to American Diabetes Association (ADA) criteria (when two or more venous plasma glucose concentrations are met or exceeded for positive diagnosis and these include: Fasting Blood Sugar (FBS)  $\geq 95$  mg/dl, BS 1 hour after 100 gr glucose  $\geq 180$  mg/dl, BS 2 hours after 100 gr glucose  $\geq 155$  mg/dl, BS 3 hours after 100 gr glucose  $\geq 140$  mg/dl). The cases with GDM were considered as case group and the women without GDM were enrolled as control group. The required data (demographic, family history, self-gestational history, anthropometric and personal habits) was recorded by trained clinical staff and collected in checklists; also, all women completed the Pittsburgh Sleep Quality Index (PSQI). The PSQI score  $< 5$  was defined as poor sleep (16).

Data analysis among 3000 women (290 cases and 2710 controls) was done by SPSS version 25.0 software. The utilized tests for comparisons across the groups were chi-square and independent samples t-test and the p values less than 0.05 were considered statistically significant. Ultimately for elimination of distorting effect of different variables, the logistic regression was used for detecting the association of sleep quality and quantity in case and control groups.

## Results

Demographic data of two groups is shown in table 1. As shown in figures 1 and 2, the gravid and parity status significantly differed across the groups (p=0.0001). As shown in figure 3, the monthly income was the same across the groups (p=0.055). As shown in figure 4, there was significant difference between quality of life in case and control groups (p=0.0001).

Different sleep quality subscales are shown in table 2 with significant difference for majority of items. As shown in table 3, the sleep duration subscales did not differ for the time of going to bed and the

**Table 1.** Demographic data across the groups

| Variables                    | GDM group      | Non diabetic group | p value |
|------------------------------|----------------|--------------------|---------|
| Mean age (years)             | 31.9           | 29.5               | 0.0001  |
| Family history of diabetes   | 91 (31.4%)     | 803 (29.6%)        | 0.536   |
| Personal history of GDM      | 44 (15.2%)     | 136 (5.0%)         | 0.0001  |
| Smoking                      | 10 (3.4%)      | 24 (0.9%)          | 0.001   |
| Undergraduate level literacy | 40 (13.8%)     | 390 (14.4%)        | 0.0001  |
| Non-Iranian patients         | 10 (3.4%)      | 10 (0.4%)          | 0.0001  |
| Mean BMI ( $kg/m^2$ )        | 27.9 $\pm$ 5.8 | 28.2 $\pm$ 4.4     | 0.413   |
| History of hypertension      | 42 (15.1%)     | 108 (4.0%)         | 0.0001  |

p- value  $\geq$  0.05 showed significant difference across two groups

**Table 2.** Sleep quality subscales across the groups

| Item                                   | GDM | Never        | Once        | Twice       | Thrice      | p value |
|--|-----|--------------|-------------|-------------|-------------|---------|
| Inability to sleep                     | No  | 860 (31.7%)  | 550 (20.3%) | 680 (25.1%) | 620 (22.9%) | 0.0001  |
|  | Yes | 80 (27.6%)   | 30 (10.3%)  | 80 (27.6%)  | 100 (34.5%) |         |
| Waking up for the need to bathe        | No  | 2410 (88.9%) | 210 (7.7%)  | 80 (3.0%)   | 10 (0.4%)   | 0.671   |
|  | Yes | 260 (89.7%)  | 20 (6.9%)   | 10 (3.4%)   | ---         |         |
| Waking up for inability to breathe     | No  | 1910 (70.5%) | 360 (13.3%) | 260 (9.6%)  | 180 (6.6%)  | 0.0001  |
|  | Yes | 170 (58.6%)  | 50 (17.2%)  | 20 (6.9%)   | 50 (17.2%)  |         |
| Waking up for coughing/snoring         | No  | 2090 (77.1%) | 400 (14.8%) | 150 (5.5%)  | 70 (2.6%)   | 0.0001  |
|  | Yes | 200 (69.0%)  | 70 (24.1%)  | 10 (3.4%)   | 10 (3.4%)   |         |
| Waking up for cold sensation           | No  | 2150 (79.3%) | 400 (14.8%) | 150 (5.5%)  | 10 (0.4%)   | 0.0001  |
|  | Yes | 190 (65.5%)  | 70 (24.1%)  | 30 (10.3%)  | ---         |         |
| Waking up for warm sensation           | No  | 1070 (39.5%) | 680 (25.1%) | 560 (20.7%) | 400 (14.8%) | 0.0001  |
|  | Yes | 110 (37.9%)  | 30 (10.3%)  | 60 (20.7%)  | 90 (31.0%)  |         |
| Waking up for nightmare                | No  | 1960 (72.3%) | 490 (18.1%) | 210 (7.7%)  | 50 (1.8%)   | 0.002   |
|  | Yes | 180 (62.1%)  | 70 (24.1%)  | 30 (10.3%)  | 10 (3.4%)   |         |
| Waking up for pain                     | No  | 1030 (38.0%) | 670 (24.7%) | 520 (19.2%) | 490 (18.1%) | 0.0001  |
|  | Yes | 80 (27.6%)   | 60 (20.7%)  | 40 (13.8%)  | 110 (37.9%) |         |
| Sedative use                           | No  | 2660 (98.2%) | 20 (0.7%)   | 10 (0.4%)   | 20 (0.7%)   | 0.0001  |
|  | Yes | 240 (82.8%)  | 40 (13.8%)  | ---         | 10 (3.4%)   |         |
| Somnolence                             | No  | 2020 (74.5%) | 500 (18.5%) | 100 (3.7%)  | 90 (3.3%)   | 0.0001  |
|  | Yes | 180 (62.1%)  | 60 (20.7%)  | 30 (10.3%)  | 20 (6.9%)   |         |
| Difficulty in doing routine activities | No  | 1110 (41.0%) | 780 (28.8%) | 580 (21.4%) | 240 (8.9%)  | 0.0001  |
|  | Yes | 100 (34.5%)  | 60 (20.7%)  | 70 (24.1%)  | 60 (20.7%)  |         |

**Table 3.** Sleep duration subscales across the groups

| Item                   | GDM | Mean | Standard deviation | p value |
|------------------------|-----|------|--------------------|---------|
| Time of going to bed   | No  | 8.05 | 10.35              | 0.831   |
|                        | Yes | 8.13 | 10.37              |         |
| Time of going to sleep | No  | 5.5  | 9.00               | 0.075   |
|                        | Yes | 4.54 | 8.22               |         |
| Wake-up time           | No  | 8.23 | 1.49               | 0.002   |
|                        | Yes | 8.42 | 1.38               |         |
| Sleep duration         | No  | 7.68 | 1.64               | 0.009   |
|                        | Yes | 8.21 | 1.48               |         |




Figure 1. Gravid status across the groups




Figure 2. Parity status across the groups




Figure 3. Monthly income across the groups




Figure 4. Sleep quality across the groups

time of sleeping ( $p>0.05$ ); but there was significant difference for wake-up time ( $p=0.002$ ) and sleep duration ( $p=0.009$ ) across the groups. The mean Pittsburgh point was  $2.31.5\pm$  and  $6.21.8\pm$  in control and cases group, respectively ( $p=0.0001$ ).

The logistic regression was predictive up to 90% ( $p=0.0001$ ) with a range of prediction between 30.1 to 64.1% (mean of 94.1%). Significant factors include previous GDM ( $p=0.006$ ), literacy ( $p=0.0001$ ), hypertension ( $p=0.032$ ), age ( $p=0.020$ ), and Pittsburgh score ( $p=0.0001$ ).

### Discussion

The results demonstrated that both quality and quantity of sleep may be related with GDM. The quality of sleep in GDM cases was worse and duration of sleep was longer. Prevalence rate of GDM is increasing worldwide with some variation according to age, ethnicity, race, body composition, and diagnostic and screening tests characteristics. There are multiple short-term and long-term complications affecting both maternal and neonatal health. Women with GDM are at increased risk for diabetes mellitus type II and cardiovascular disorders. Children of diabetic mothers will have increased risk for childhood obesity, abnormal glucose level, and early-onset cardiovascular diseases. GDM is a carbohydrate intolerance status during pregnancy either treated with or without insulin. More than half of GDM cases would develop overt diabetes mellitus till the next 20 years. Hence, recognition of related factors would help in better prevention of GDM. Sleep disorders especially elongated sleep, poor-quality sleep, and sleep-breathing disorders are common during pregnancy which may be connected with hormonal changes, anxiety, and somatic problems.

In non-pregnant population, it has been shown by Chaput *et al* (7) that increased and decreased night-time sleep would increase the incidence rate of diabetes mellitus. Wang *et al* (10) reported that poor-quality sleep and sleep duration longer than nine hours versus 7 to 9 hours are accompanied with higher incidence rate of GDM. Similarly, in our study, the sleep quality was worse in mothers with GDM and also the longer duration was related to GDM. In some studies, the economic and social status, income levels, and literacy were separately related to sleep

disorders and occurrence of GDM (16,17).

The study by Saadati *et al* (18) reported that poor-sleep quality was related to mild preeclampsia and GDM. Other studies (19-22) showed that short duration of sleep was related to GDM. In this study, there was no significant difference between family history among healthy and GDM mothers. It is recommended that in future studies, the second-degree relatives be evaluated as well.

Sleeping difficulties, difficulty breathing or shortness of breath, coughing or snoring, nightmare, and pain are more common among diabetic cases. Also, the use of sedatives was more common among GDM mothers. The mean Pittsburgh score was 6.2 that showed poor sleep quality similar to other studies (18,19,23). Regarding the obtained association between quality and quantity of sleep with GDM, some training courses and guidance would result in improved sleep pattern

and subsequent decrease in GDM incidence rate.

Totally, according to the obtained results, it may be concluded that quality and quantity of sleep is associated with the incidence of gestational diabetes and diabetic mothers have worse sleep quality and longer sleep times than non-diabetic mothers. However, further studies with larger sample size are required to attain more definite results.

## Conclusion

This study showed significant association between quality and quantity of sleep and GDM in pregnancy, although studies with larger sample size are required and the priority of sleep disorders in occurrence of GDM needs to be more investigated. That may result in early detection of GDM in these patients or can lead to decrease of GDM prevalence by treating sleep disorders during pregnancy.

## References

- Hillier TA, Pedula KL, Schmidt MM, Mullen JA, Charles MA, Pettitt DJ. Childhood obesity and metabolic imprinting: the ongoing effects of maternal hyperglycemia. *Diabetes Care* 2007;30(9):2287-92.
- Silverman BL, Metzger BE, Cho NH, Loeb CA. Impaired glucose tolerance in adolescent offspring of diabetic mothers. Relationship to fetal hyperinsulinism. *Diabetes Care* 1995;18(5):611-7.
- Okun ML, Luther JF, Wisniewski SR, Sit D, Prairie BA, Wisner KL. Disturbed sleep, a novel risk factor for preterm birth. *J Women's Health (Larchmt)* 2012;21(1):54-60.
- Okun ML, Roberts JM, Marsland AL, Hall M. How disturbed sleep may be a risk factor for adverse pregnancy outcomes. *Obstet Gynecol Surv* 2009;64(4):273-80.
- Reutrakul S, Zaidi N, Wroblewski K, Kay HH, Ismail M, Ehrmann DA, et al. Sleep disturbances and their relationship to glucose tolerance in pregnancy. *Diabetes Care* 2011;34(11):2454-7.
- Ayas NT, White DP, Al-Delaimy WK, Manson JE, Stampfer MJ, Speizer FE, et al. A prospective study of self-reported sleep duration and incident diabetes in women. *Diabetes Care* 2003;26(2):380-4.
- Chaput JP, Despres JP, Bouchard C, Tremblay A. Association of sleep duration with type 2 diabetes and impaired glucose tolerance. *Diabetologia* 2007;50(11):2298-304.
- Gangwisch JE, Heymsfield SB, Boden-Albala B, Buijs RM, Kreier F, Pickering TG, et al. Sleep duration as a risk factor for diabetes incidence in a large US sample. *Sleep* 2007;30(12): 1667-73.
- Luque-Fernandez MA, Bain PA, Gelaye B, Redline S, Williams MA. Sleep-disordered breathing and gestational diabetes mellitus: a meta-analysis of 9,795 participants enrolled in epidemiological observational studies. *Diabetes Care* 2013;36(10):3353-60.
- Wang H, Leng J, Li W, Wang L, Zhang C, Li W, et al. Sleep duration and quality, and risk of gestational diabetes mellitus in pregnant Chinese women. *Diab Med* 2017;34(1):44-50.
- Lzci-Balserak B, Pien GW. The relationship and potential mechanistic pathways between sleep disturbances and maternal hyperglycemia. *Curr Diab Rep* 2014;14(2):459.

12. O’Keeffe M, St-Onge MP. Sleep duration and disorders in pregnancy; implications for glucose metabolism and pregnancy outcomes. *Int J Obes (Lond)* 2013;37(6):765-70.
13. Herring SJ, Nelson DB, Pien GW, Homko C, Goetzl LM, Davey A, et al. Objectively- measured sleep duration and hyperglycemia in pregnancy. *Sleep Med* 2014;15(1):51-5.
14. Kita T, Yoshioka E, Satoh H, Saijo Y, Kawaharada M, Okada, E et al. Short sleep duration and poor sleep quality increase the risk of diabetes in Japanese workers with no family history of diabetes. *Diabetes Care* 2012;35(2):313-8.
15. Mallon L, Broman JE, Hetta J. High incidence of diabetes in men with sleep complaints or short sleep duration: a 12-year follow-up study of a middle-aged population. *Diabetes Care* 2005;28(11):2762-7.
16. Bo S, Menato G, Bardelli C, Lezo A, Signorile A, Repeni E, et al. Low socioeconomic status as a risk factor for gestational diabetes. *Diabetes Metab* 2002;28(2):139-40.
17. Signal TL, Paine SJ, Sweeney B, Priston M, Muller D, Smith A, et al. Prevalence of abnormal sleep duration and excessive daytime sleepiness in pregnancy and the role of socio-demographic factors: comparing pregnant women with women in the general population. *Sleep Med* 2014;15(12):1477-83.
18. Saadati F, Sehhatiei Shafaei F, Mirghafourvand M. Sleep quality and its relationship with quality of life among high-risk pregnant women (gestational diabetes and hypertension). *J Matern Fetal Neonatal Med* 2018;31(2):150-7.
19. Hayase M, Shimada M, Seki H. Sleep quality and stress in women with pregnancy-induced hypertension and gestational diabetes mellitus. *Women Birth* 2014;27(3):190-5.
20. Facco FL, Grobman WA, Kramer J, Ho KH, Zee PC. Self-reported short sleep duration and frequent snoring in pregnancy: impact on glucose metabolism. *Am J Obstet Gynecol* 2010;203(2):142.e1-142.e5.
21. Facco FL, Grobman WA, Reid KJ, Parker CB, Hunter SM, Silver RM, et al. Objectively measured short sleep duration and later sleep midpoint in pregnancy are associated with a higher risk of gestational diabetes. *Am J Obstet Gynecol* 2017;217(4):447.e1-447.e13.
22. Ferrari U, Kunzel H, Trondle K, Rottenkolber M, Kohn D, Fugmann M, et al. Poor sleep quality is associated with impaired glucose tolerance in women after gestational diabetes. *J Psych Res* 2016;65:166-71.
23. Cai S, Tan S, Gluckman D, Godfrey KM, Saw SM, Teoh OH, et al. Sleep quality and nocturnal sleep duration in pregnancy and risk of gestational diabetes mellitus. *Sleep* 2017;40(2):201-9.