

Involvement of NMDA Receptor and Nitric Oxide Pathway in the Anticonvulsant Effect of Genistein in Ovariectomized Mice

Mohammad Algazo^{1,2}, Nastaran Rahimi^{1,2}, Saeed Amiri Gheshlaghi^{1,2}, Hussein Alshaib^{1,2}, Reza Fahimi^{1,2}, Ahmad Reza Dehpour^{1,2*}

1. Experimental Medicine Research Center, Tehran University of Medical Sciences, Tehran, Iran

2. Department of Pharmacology, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Abstract

Background: Genistein is an isoflavonoid of soy that has been stated to show neuroprotection effect in some central nervous system disorders such as seizures and status epilepticus. From our previous study, we found that genistein showed an anticonvulsant effect through estrogenic and serotonergic receptors in Ovariectomized (OVX) mice. N-methyl d-aspartate (NMDA) receptors and Nitric Oxide (NO) pathway have been documented to be involved in genistein effects in rat's hippocampus ischemic/reperfusion and Alzheimer's disease. Accordingly, we aimed to evaluate the possible involvement of NMDA receptors and nitric oxide pathway in anticonvulsant effect of genistein on pentylenetetrazole (PTZ) induced seizures in OVX-mice.

Methods: NMRI female mice weighing 23-30 gr were undergone bilateral ovariectomy. Seizure susceptibility was studied by PTZ-induced seizures model after 14 days of recovery.

Results: Genistein (10 mg/kg, i.p.) injection elevated the seizure threshold in OVX mice. The effect of a sub-effective dose of genistein (5 mg/kg, i.p.) was potentiated by L-NAME (nonspecific inhibitor of nitric oxide synthase (NOS); 10 mg/kg, i.p.) and 7-nitroindazole (specific inhibitor of neural NOS; 25 mg/kg, i.p.), while, L-arginine (the nitric oxide precursor, 30 mg/kg, i.p.) blocked the anticonvulsant activity of genistein (10 mg/kg). Acute injection of ketamine (0.5 mg/kg, i.p.) and MK-801 (0.05 mg/kg) as NMDA receptor antagonists before sub-effective dose of genistein (5 mg/kg, i.p.) showed a significant increase in seizure threshold among OVX mice.

Conclusion: In summary, our results demonstrated that NMDA receptors and neuronal nitric oxide synthase might be associated with an anticonvulsant effect of genistein after ovariectomy in mice.

Keywords: Genistein, Mice, Nitric Oxide, NMDA Receptor, Ovariectomy, Seizure Threshold

* Corresponding author

Ahmad Reza Dehpour, PhD

Experimental Medicine Research Center,
Tehran University of Medical Sciences,
Tehran, Iran

Tel: +98 21 88973652

Fax: +98 21 66402569

Email: dehpoura@sina.tums.ac.ir

Received: 17 Jan 2019

Accepted: 20 Mar 2019

Citation to this article:

Algazo M, Rahimi N, Amiri Gheshlaghi S, Alshaib H, Fahimi R, Dehpour AR. Involvement of NMDA Receptor and Nitric Oxide Pathway in the Anticonvulsant Effect of Genistein in Ovariectomized Mice. *J Iran Med Council.* 2019;1(3): 133-139.

Introduction

Genistein is the soy isoflavone¹ which can pass across the blood-brain barrier like estradiol². Several studies reported that genistein plays a role as a powerful neuroprotective in central nervous system³⁻⁶. Also, it could bind to estrogen receptors and has been reported to act like an agonist of both α and β estrogen receptors from different studies on cell culture, estrogen receptor, reporter-genes and animal models⁷. In our previous experiment, genistein was able to significantly increase the seizure threshold induced by pentylenetetrazole in Ovariectomized (OVX) mice. This anticonvulsant impact of genistein was prohibited by selective antagonist of estrogen receptor⁸. Also, genistein treatment has neuroprotective effects on status epilepticus-induced hippocampal damage in OVX female rats and preserved the inhibition of the dentate gyrus^{2,7}. Menze *et al* in 2015 revealed a neuroprotective and memory enhancing effects of genistein in a rat model of Huntington's disease. Also they suggest these impacts of genistein was mediated by its antioxidant, anti-inflammatory and cholinesterase inhibitory activities³. Moreover, it was shown that genistein inhibited glutamate-induced apoptosis in mouse primary neuronal cell cultures⁴.

Genistein is known as a powerful phytoestrogen. Phytoestrogens can obviously increase the expression of Nitric Oxide Synthase (NOS) in hippocampus playing a protecting role against neurodegenerative disease⁹. Pretreatment with genistein and 17 β -estradiol in Alzheimer's disease decreased the expression of inducible Nitric Oxide Synthase (iNOS) in cultured astrocytes¹⁰. Furthermore, our previous study has shown nonselective nitric oxide synthase inhibitor blocked the anticonvulsant activity of oleuropein, which is another well-known phytoestrogen¹¹.

In addition, N-methyl-D-aspartate (NMDA) receptors have received very much attention because of their involvement in neuronal development, neurodegenerative diseases, and neuronal excitotoxicity¹²⁻¹⁴. Genistein significantly inhibited glutamate-induced apoptosis and the effect of this isoflavone was most prominent in the hippocampal cells⁵. Earlier studies revealed that genistein not only significantly depressed ischemia/reperfusion-related release of glutamate among rat's cerebral cortex, but also reduced the influx of Ca²⁺ after cerebral ischemia through inhibiting tyrosine-phosphorylation of NMDA receptors¹⁵.

In present study, we intended to assess the possible neuro-protective mechanisms of genistein on the seizures threshold through NMDA receptors/NO pathway.

Materials and Methods

Animals

Mice (23-30 g) were obtained from Department of Pharmacology of Tehran University of Medical Sciences. Animals were accommodated under standard laboratory conditions including temperature (23 \pm °C), a 12-hr-dark/12-hr-light cycle in polycarbonate cages with free access to standard animal chow and water. The study was conducted in accordance with the Guidelines for the Care and Use of Laboratory Animals published by the National Institutes of Health (NIH publication NO. 85-23; revised 1985).

Materials

The following materials were used: Genistein, pentylenetetrazole (PTZ), MK-801 (an uncompetitive antagonist of the N-Methyl-D-aspartate (NMDA) receptor, glutamate receptor), L-NG-Nitro-L-arginine methyl ester hydrochloride (L-NAME, a non-selective inhibitor of NOS), L-arginine (a NO precursor) that all were purchased from Sigma (St. Louis, MO, USA). Ketamine hydrochloride (a nonbarbiturate anesthetic) was obtained from Rotexmedica (Trittau, Germany) and xylazine hydrochloride purchased from Alfasan (Woerden, The Netherlands).

The chemicals (L-NAME, L-arginine and ketamine) were dissolved/diluted in sterile isotonic saline solution; Genistein and Mk-01 dissolved in dimethyl sulfoxide (DMSO)/PBS at a ratio of 1:6. All chemicals were administered intraperitoneally (i.p.) in a volume of 10 ml/kg for the mice body weight, however pentylenetetrazole (PTZ; 0.5%) was excepted and injected intravenously (i.v.). Appropriate vehicle controls were prepared for each experiment. At least eight animals were used for each treatment group and each mouse was tested only once.

All behavioral tests were performed after a recovery period of 14 days of induction for ovariectomized¹⁶⁻¹⁸.

Determination of seizure susceptibility

A more sensitive method of i.v. administration of PTZ that allows better detection modulatory effects on convulsive tendency was used to assess the seizure

susceptibility among mice. The threshold of PTZ was determined by inserting a 30-gauge butterfly needle into the tail vein of mice while they were restrained in a mouse restrainer. The needle was then secured to the tail with a narrow piece of adhesive tape, with the mouse moving freely. PTZ (0.5%) was infused at the constant rate of 1 ml/min using an infusion pump. Infusion was halted when forelimb clonus followed by a full clonus of the whole body was observed and the dose of PTZ administered (mg/kg of mice weight) was measured as an index of clonic seizure threshold knowing that the seizure threshold is dependent on PTZ dosage and is also time-related¹⁹.

Experiments

Intraperitoneal (i.p.) administration of genistein (10 mg/kg) was performed in three different time intervals (15, 30, 45 min). Then different doses of genistein (5, 10, 30 mg/kg, i.p.) was injected 30 min prior to induction of seizure. Doses and time points were determined based on a pilot experiment. Control animals received DMSO 1%. Administration of L-NAME (10 mg/kg, i.p.) and 7-nitroindazole (25 mg/kg, i.p.) was carried out 15 min before sub-effective dose of genistein (5 mg/kg). L-arginine (60 mg/kg, i.p.) was injected to mice 15 min before vehicle and genistein (10 mg/kg) groups. MK-801 (0.05 mg/kg, i.p.) and ketamine 0.5 mg/kg (i.p.) were administered in mice 15 min prior to injection of vehicle- and genistein- treated mice. In the last experiment, diazepam (0.05 mg/kg, i.p.) was used as a positive control.

Statistical analysis

Data are presented as mean \pm SEM of PTZ minimal dose (mg/kg) and analyzed using the SPSS statistical software package (Version 22). The data of PTZ-induced seizure thresholds were analyzed by one or two way analysis of variance (ANOVAs) followed by post hoc Tukey's tests in two group comparison and multiple group comparison, respectively. The tests of homogeneity of variance were used to ensure normal distribution of the data. A p-value less than 0.05 were defined as statistically significant level.

Results

The effect of time and different doses of genistein on

the seizure threshold

Figure 1 shows the time of the genistein effect on the seizure threshold. As shown, intraperitoneal administration of genistein (10 mg/kg) did not affect the PTZ-induced seizure threshold at 15 and 45 min, whereas it significantly reduced the seizure threshold 30 min after administration ($p < 0.001$).

Figure 2 illustrates the effect of acute intraperitoneal administration of genistein (5, 10, and 30 mg/kg, i.p.) on the seizure threshold, when it was administered 30 min prior to PTZ. Genistein (10 mg/kg) significantly increased the PTZ-induced seizure threshold compared with vehicle-treated control animals ($p < 0.001$).




Figure 1. Effect of genistein (10 mg/kg, i.p.) on PTZ-induced clonic seizure threshold in mice. As shown, genistein was administered 15, 30 and 45 min before PTZ. The effects were compared with a control group at the same time. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice. *** $p < 0.001$ compared with corresponding vehicle group.




Figure 2. The effect of different doses of genistein on the PTZ induced seizure threshold in mice. Injection of genistein was performed 30 min before induced seizure and diazepam (0.05 mg/kg) was injected as a positive control. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice. * $p < 0.05$, *** $p < 0.001$ compared with vehicle group.

Impacts of nitric oxide pathway on the anticonvulsive effect of genistein in PTZ-induced seizures in mice

Figures 3 and 4 shows the effects of sub-effective dose of L-NAME and 7-nitroindazole pretreatment on

sub-effective dose of genistein on seizure threshold in PTZ-induced seizures in mice. Administration of L-NAME (10 mg/kg, i.p.) and 7-nitroindazole (25 mg/kg, i.p.) by their own had no effect on seizure threshold ($p>0.05$) in comparison with vehicle treated controls. While, administration of both of them 15 min before sub-effect dose of genistein (5 mg/kg, i.p.) significantly increased the anti-convulsant effect of genistein ($p<0.001$).

Figure 5 shows the L-arginine pretreatment effect on the anticonvulsant activity of genistein against PTZ-induced seizures. As shown in this figure, L-arginine pretreatment significantly reversed the anticonvulsant effect of genistein ($p<0.001$). Administration of L-arginine by its own had no effect on seizure threshold in comparison with saline treated controls.

Impacts of N-methyl-D-aspartate (NMDA) receptors on the convulsive effect of genistein in PTZ-induced seizures in mice

As shown in figure 6, administration of ketamine as a non-competitive antagonist of NMDA receptor (0.5 mg/kg, i.p.) did not alter the seizure threshold induced by PTZ in mice, while acute administration of ketamine (0.5 mg/kg, i.p.) 15 min before genistein (5 mg/kg, i.p.) significantly enhanced the anticonvulsant effect of genistein ($p<0.001$).

Furthermore, acute injection of MK-801 (a potential noncompetitive antagonist of NMDA receptor) 15 min




Figure 3. this chart shows the sub-effect dose of L-NAME pretreatment on sub-effect dose of genistein on PTZ-induced seizures. Administration of L-NAME by its own had no effect on seizure threshold in comparison with control group, while administration of L-NAME before sub-effective dose of genistein augmented the anticonvulsant effect of genistein. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice.

*** $p<0.001$ compared with corresponding vehicle group; @@@ $p<0.001$ compared with genistein (5 mg/kg) treated group.




Figure 4. Injection of 7-nitroindazole before vehicle (solvent) had no effect on seizure threshold in comparison with vehicle treated, mice however co-administration of 7-nitroindazole with genistein increased the anticonvulsant effect of genistein. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice.

*** $p<0.001$ compared with corresponding vehicle group; @@@ $p<0.001$ compared with genistein (5 mg/kg) treated group.




Figure 5. This figure shows the effect of L-arginine pretreatment on the anticonvulsant effect of genistein in PTZ-induced seizures. As shown, L-arginine pretreatment significantly reversed the anticonvulsant effect of genistein. Administration of L-arginine by its own did not alter the seizure threshold in comparison with saline treated controls. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice. *** $p<0.001$ compared with corresponding vehicle group; #### $p<0.001$ compared with genistein (10 mg/kg) treated group.




Figure 6. Administration of ketamine significantly decreased the seizure threshold in comparison with genistein-treated controls. Furthermore, genistein pretreatment led to an increase in seizure threshold when compared with control group. Ketamine by its own had no effect on seizure threshold. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice. *** $p<0.001$ compared with corresponding vehicle group; @@@ $p<0.001$ compared with genistein (5 mg/kg) treated group.

before genistein (10 mg/kg, i.p.) significantly increased the anticonvulsant effect of genistein (5 mg/kg, i.p.) ($p < 0.001$; Figure 7), whereas, MK-801 by its own had no significant effect on seizure threshold.




Figure 7. This figure illustrates the effect of MK-801 (a potent N-methyl-D-aspartate antagonist) on the PTZ-induced seizures. MK-801 pretreatment increased the anticonvulsant effect of genistein. MK-801 by its own had no effect on seizure threshold. Data are expressed as means \pm SEM of seizure threshold in at least 8 mice. *** $p < 0.001$ compared with corresponding vehicle group; @@@ $p < 0.001$ compared with genistein (5 mg/kg) treated group.

Discussion

We found that genistein have anticonvulsant characteristics in PTZ-induced seizure model in ovariectomized mice. L-arginine inhibited genistein effect on seizures threshold while L-NAME and 7-nitroindazole with sub-effective dose of genistein exacerbated the anticonvulsant effect of sub-effective dose of genistein. In our study, blocking of NMDA receptor with either ketamine or MK-80 led to inhibition of anticonvulsant effect of genistein. We, then, propose that genistein can modulate the anticonvulsant effect throughout NMDA receptors and nitrenergic pathway in ovariectomized mice.

Recently Westmark *et al* investigated the effect of soy-diet on seizure in neurologic disorders such as Alzheimer's disease, Down syndrome and Fragile X Syndrome (FXS) in audiogenic seizure mouse models. Surprisingly they found that daidzein exacerbated seizure events in mice which suffered from neurologic disorders²⁰. In our previous study, we found that estrogenic receptors are involved in genistein anticonvulsant effect among mice after ovariectomy⁸.

Nitric oxide has been reported to act as a neuromodulator that exhibits dual pro-convulsive^{19,21} and anticonvulsive^{11,22} properties in PTZ model. In the present study we found that L-arginine inhibited genistein effect on seizure

while L-NAME with sub-effective dose of genistein exacerbated the anticonvulsant effect of non-effective dose of genistein. Our results show that NO modulation is involved in the anticonvulsant effect of genistein so these results come in line with previous studies that have shown that nitric oxide system modulation was one of the most used mechanisms that genistein delivers its action through them^{5,12,23}. Valsecchi *et al* concluded that NOS inhibition could be one of the multiple mechanisms that genistein relieved neuropathic pain, with both peripheral and CNS origins²³. Liu *et al* demonstrated that genistein had direct non genomic effects on NOS activity on vascular endothelial cells leading to NOS activation that these effects were mediated by protein kinase activity and were unrelated to an estrogenic effect³. Gingerich *et al* revealed that genistein decreased the number of nNOS-positive neurons in PVN through activation of beta estrogen receptors by genistein²⁴. Estrogen has an effect on NMDA in brain²⁵. NMDA/NO cascade is a well-known pathway in brain where NMDA mediates seizure events and behavioral changes by activating NO system²⁶⁻²⁹.

In our study, blocking NMDA receptor by either ketamine or MK-801 led to inhibition of the anticonvulsant effect of genistein. These findings come in line with previous studies that showed NMDA receptor was implicated in mediating some of genistein effects. It has found that genistein inhibited NMDA-dependent cGMP production with high potencies³⁰⁻³². Another study concluded estrogen receptor-mediated stimulation of the nNOS/PSD-95/NMDA receptor was likely to be a critical component of the signaling process by which estradiol facilitated coupling of glutamatergic fluxes for NO production in neurons³³. Taken together, to explain our findings we suggest that mechanisms which underlines the anticonvulsant effect of genistein at least is partly carried out by modulating estrogen receptor/NMDA/ NO cGMP pathway which at the end leads to regulation of NO synthesis in brain.

Conclusion

In conclusion, the current findings underline that genistein had anticonvulsant activity in PTZ-induced seizures. This phenomenon could be modulated by NMDA receptor antagonists and nitric oxide synthase mediators which suggests the involvement of NMDA/ nitrenergic pathways within the anticonvulsive impacts

of genistein in the PTZ model of clonic seizures among ovariectomized mice.

Medicine Research Center, Tehran University of Medical Sciences (grant No. 91-02-158-18207).

Acknowledgments

This study was supported by a grant from Experimental

Conflict of Interest

Authors declare that there is no conflict of interest.

References

1. Naaz A, Yellayi S, Zakroczymski MA, Bunick D, Doerge DR, Lubahn DB, et al. The soy isoflavone genistein decreases adipose deposition in mice. *Endocrinology* 2003;144(8):3315-20.
2. Melcangi RC, Panzica GC. Neuroactive steroids: old players in a new game. *Neuroscience* 2006;138(3):733-9.
3. Menze ET, Esmat A, Tadros MG, Abdel-Naim AB, Khalifa AE. Genistein improves 3-NPA-induced memory impairment in ovariectomized rats: impact of its antioxidant, anti-inflammatory and acetylcholinesterase modulatory properties. *PloS One* 2015;10(2):e0117223.
4. Kajta M, Domin H, Gryniewicz G, Lason W. Genistein inhibits glutamate-induced apoptotic processes in primary neuronal cell cultures: An involvement of aryl hydrocarbon receptor and estrogen receptor/glycogen synthase kinase-3 β intracellular signaling pathway. *Neuroscience* 2007;145(2):592-04.
5. Lund TD, West TW, Tian LY, Bu LH, Simmons DL, Setchell KD, et al. Visual spatial memory is enhanced in female rats (but inhibited in males) by dietary soy phytoestrogens. *BMC Neurosci* 2001;2(1):20.
6. Coldham NG, Sauer MJ. Pharmacokinetics of [14 C] genistein in the rat: gender-related differences, potential mechanisms of biological action, and implications for human health. *Toxicol Appl Pharmacol* 2000;164(2):206-15.
7. Bhat RV, Budd Haeberlein SL, Avila J. Glycogen synthase kinase 3: a drug target for CNS therapies. *J Neurochem* 2004;89(6):1313-7.
8. Gheshlaghi SA, Jafari RM, Algazo M, Rahimi N, Alshaib H, Dehpour AR. Genistein modulation of seizure: involvement of estrogen and serotonin receptors. *J Natl Med* 2017;71(3):537-44.
9. Huang TC, Lu KT, Wo YY, Wu YJ, Yang YL. Resveratrol protects rats from A β -induced neurotoxicity by the reduction of iNOS expression and lipid peroxidation. *PloS One* 2011;6(12):e29102.
10. Guo K, Xu J, Yuan Q, Huang R. Effects of phytoestrogens on NOS positive neurons in hippocampus and on cognitive functions of ovariectomized rats. *Chinese J Anatomy* 2003;27(4):404-7.
11. Rahimi N, Delfan B, Motamed-Gorji N, Dehpour AR. Effects of oleuropein on pentylenetetrazol-induced seizures in mice: involvement of opioidergic and nitrenergic systems. *J Nat Med* 2017;71(2):389-96.
12. Marquina M, España A, Fernández-Galar M, López-Zabalza MJ. The role of nitric oxide synthases in pemphigus vulgaris in a mouse model. *British J Dermatol* 2008;159(1):68-76.
13. Dingledine R, Borges K, Bowie D, Traynelis SF. The glutamate receptor ion channels. *Pharmacol Rev* 1999;51(1):7-62.
14. Cull-Candy S, Brickley S, Farrant M. NMDA receptor subunits: diversity, development and disease. *Curr Opin Neurobiol* 2001;11(3):327-35.
15. Candelario-Jalil E, Muñoz E, Fiebich BL. Detrimental effects of tropisetron on permanent ischemic stroke in the rat. *BMC Neurosci* 2008;9(1):19.
16. Kalbasi Anaraki D, Sianati S, Sadeghi M, Ghasemi M, Paydar MJ, Ejtemaei Mehr S, et al. Modulation by female sex hormones of the cannabinoid-induced catalepsy and analgesia in ovariectomized mice. *Eur J Pharmacol*

2008;586(1):189-96.

17. Sadeghi M, Sianati S, Anaraki DK, Ghasemi M, Paydar MJ, Sharif B, et al. Study of morphine-induced dependence in gonadectomized male and female mice. *Pharmacol Biochem Behav* 2008;91(4):604-9.

18. Heydarpour P, Salehi-Sadaghiani M, Javadi-Paydar M, Rahimian R, Fakhfour G, Khosravi M, et al. Estradiol reduces depressive-like behavior through inhibiting nitric oxide/cyclic GMP pathway in ovariectomized mice. *Horm Behav* 2013;63(2):361-9.

19. Javadian N, Rahimi N, Javadi-Paydar M, Doustimotlagh AH, Dehpour AR. The modulatory effect of nitric oxide in pro-and anti-convulsive effects of vasopressin in PTZ-induced seizures threshold in mice. *Epilepsy Res* 2016;126:134-40.

20. Westmark CJ, Westmark PR, Malter JS. Soy-based diet exacerbates seizures in mouse models of neurological disease. *J Alzheimers Dis* 2013;33(3):797-05.

21. Rahimi N, Sadeghzadeh M, Javadi-Paydar M, Heidary MR, Jazaeri F, Dehpour AR. Effects of D-penicillamine on pentylentetrazole-induced seizures in mice: Involvement of nitric oxide/NMDA pathways. *Epilepsy Behav* 2014;39:42-7.

22. Hassanipour M, Shirzadian A, Boojar MM, Abkhoo A, Abkhoo A, Delazar S, et al. Possible involvement of nitrenergic and opioidergic systems in the modulatory effect of acute chloroquine treatment on pentylentetrazol induced convulsions in mice. *Brain Res Bull* 2016;121:124-30.

23. Valsecchi AE, Franchi S, Panerai AE, Sacerdote P, Trovato AE, Colleoni M. Genistein, a natural phytoestrogen from soy, relieves neuropathic pain following chronic constriction sciatic nerve injury in mice: anti-inflammatory and antioxidant activity. *J Neurochem* 2008;107(1):230-40.

24. Gingerich S, Krukoff TL. Estrogen modulates endothelial and neuronal nitric oxide synthase expression via an estrogen receptor β -dependent mechanism in hypothalamic slice cultures. *Endocrinology* 2005;146(7):2933-41.

25. Hoffman GE, Moore N, Fiskum G, Murphy AZ. Ovarian steroid modulation of seizure severity and hippocampal cell death after kainic acid treatment. *Exp Neurol* 2003;182(1):124-34.

26. Hilgier W, Oja SS, Saransaari P, Albrecht J. A novel glycine site-specific N-methyl-D-aspartate receptor antagonist prevents activation of the NMDA/NO/CGMP pathway by ammonia. *Brain Res* 2004;1015(1):186-8.

27. Montague PR, Gancayco CD, Winn MJ, Marchase RB, Friedlander MJ. Role of NO production in NMDA receptor-mediated neurotransmitter release in cerebral cortex. *Science* 1994;263(5149):973-6.

28. Garthwaite J, Garthwaite G, Palmer RM, Moncada S. NMDA receptor activation induces nitric oxide synthesis from arginine in rat brain slices. *Eur J Pharmacol* 1989;172(4-5):413-6.

29. Piedrafita B, Cauli O, Montoliu C, Felipe V. The function of the glutamate–nitric oxide–cGMP pathway in brain in vivo and learning ability decrease in parallel in mature compared with young rats. *Learn Mem* 2007;14(4):254-8.

30. Rodriguez J, Quignard JF, Fagni L, Lafon-Cazal M, Bockaert J. Blockade of nitric oxide synthesis by tyrosine kinase inhibitors in neurones. *Neuropharmacology* 1994;33(11):1267-74.

31. Ceccatelli S, Grandison L, Scott RE, Pfaff DW, Kow L-M. Estradiol regulation of nitric oxide synthase mRNAs in rat hypothalamus. *Neuroendocrinology* 1996;64(5):357-63.

32. McNeill AM, Kim N, Duckles SP, Krause DN, Kontos HA. Chronic estrogen treatment increases levels of endothelial nitric oxide synthase protein in rat cerebral microvessels. *Stroke* 1999;30(10):2186-90.

33. D'Anglemont de Tassigny X, Campagne C, Steculorum S, Prevot V. Estradiol induces physical association of neuronal nitric oxide synthase with NMDA receptor and promotes nitric oxide formation via estrogen receptor activation in primary neuronal cultures. *J Neurochem* 2009;109(1):214-24.